

Manipulating the scattering length of a Bose-Einstein condensate in an amplitude - modulated optical lattice.

G.M.Genkin*.

Department of Physics and Astronomy, Northwestern University , Evanston , Illinois , 60208 .

Abstract

The scattering length in a BEC confined in an amplitude - modulated optical lattice can be manipulated by the modulation strength. Two standing laser waves of main and sideband frequencies of an optical lattice induce a Raman transition; due to resonant Raman driving the effective scattering length depends on the modulation strength and the detuning from resonance and can be tuned to a given value.

* Electronic address: gena@pluto.phys.nwu.edu.

03.75.Lm, 05.30.Jp.

The recent experimental observation of Bose-Einstein condensation in a dilute gas of ultracold trapped atoms [1-3] has generated much interest in studying the properties of a Bose-Einstein condensate (BEC) and manipulating such coherent matter by external electromagnetic fields. The experimental observation of Bose-Einstein condensation in a dilute gas of ultracold trapped atoms [1-3] has generated much interest in manipulating such coherent matter by external electromagnetic fields. Recently atoms have been confined in optical potentials created by standing light waves [4, 5], with the wavelength of the optical potential much smaller than the dimensions of the trap. The unique prospects of this new class of system (the correlated bosons on a lattice) became evident. The possibility of creating optical lattice in trapped Bose - condensed gases has provided an opportunity to study superfluids in novel situations. The presence of the lattice leads to a variety of solid - state effects. For example, a transition to a Mott insulator becomes possible when a superfluid like a BEC is placed in a periodic potential [4]; the oscillation frequency of the center - of - mass motion of the condensate is reduced [6] as a result of the enhanced effective mass of the atoms tunneling between potential wells. By accelerating the optical lattice [7] Bloch oscillations of the condensate have been observed, and reducing the amplitude of the lattice optical potential leads to a breakdown of these oscillations as a result of Landau - Zener tunneling between bands.

Many physical properties of a dilute gas are determined by the atom - atom interaction. In a dilute and cold gas only binary collisions at low energy are relevant and these collisions are characterized by a single parameter, the s - wave scattering length. The confinement of a gas in one or more spatial dimensions can strongly modify the collisional properties (two - body scattering in an optical lattice) of atoms [8 - 12]. In Ref.[12] was shown that in an 3D optical lattice a dilute gas is described by the effective atomic scattering length a_{eff} , modified and tunable by the lattice parameters,

$$a_{eff} = \frac{\pi \ln 2}{4} a \frac{d}{a + l_*} \frac{\hbar}{m \omega l_0^2},$$

$$l_* = \frac{\ln 2}{4} l_0 \sqrt{D_0}, \quad (1)$$

where a is the free - space scattering length, the optical lattice potential is determined by a V_0 and lattice spacing d , and l_0 is the size of the ground state wave function in an individual well of an optical lattice. The wave functions of the relative motion of two particles of mass m are characterized by the quasimomentum q , the band index s , and the energy $\varepsilon_{sq} = \varepsilon_s - t_s \cos(qd)$, where $t_s = \sqrt{D_s} \hbar \omega / \pi$, $D_s \ll 1$ is WKB tunneling exponent between the neighboring wells, $\omega^2 = V_0 / (md^2)$, ε_s are the energies of states s in isolated wells. The effective scattering length a_{eff} is modified by the optical lattice if the tunneling between the neighboring lattice sites is sufficiently small and the new length scale l_* contains the tunneling amplitude (for the band index $s = 0$ the amplitude D_0) and describes the influence of particle states extending over several lattice sites on the two - body scattering in an optical lattice. In Ref. [12] was shown that in the case of attractive interaction, for the free - space length $a < 0$, the effective scattering length exhibits resonant behavior and leads to a geometric resonance at $l_* = |a|$, and for $l_* > |a|$ the effective scattering becomes repulsive $a_{eff} > 0$.

The goal of this paper is manipulating of the effective scattering length by a time - dependent action of the lattice parameters. We consider a manipulation of the scattering length in a BEC in an amplitude - modulated optical lattice. The such optical lattice can be created by interfering pair of amplitude - modulated beams. In general, there are the two generic types (a resonant and a nonresonant cases) of the manipulation. The effect of the modulation is to modulate the optical lattice potential so that it becomes a time - dependent optical lattice potential which the depth of the optical potential is

$$V_0(t) = V_0(1 + \nu \cos \omega_M t), \nu < 1, \quad (2)$$

where ν is the modulation strength, ω_M is the modulation frequency. If the period of modulation $2\pi/\omega_M$ is larger the nonequilibrium dynamic time τ_{noneq} , then the time behavior of a BEC in an amplitude - modulated optical lattice is quasistatic. The nonequilibrium dynamic time τ_{noneq} for trapped bosonic atoms in an optical lattice potential was considered in Ref.[11] and was shown that the characteristic time of the dynamical restoration of the

phase coherence is a Josephson time $\hbar/J$.

A resonant case.

One is a resonant mechanism which operates when the modulation frequency ω_M is close to the excited internal state of a BEC. A resonant mechanism is determined by change in the population of the ground state due to a resonant driving. In a BEC in a periodic potential the characteristic energetic scale of the band structure [14,15] is the recoil energy E_R . And, usually, E_R exceeds J , and the quasistatic behavior is not valid. However, in this case is more convenient to use a picture that an amplitude - modulated field can be presented as a set of monochromatic fields (main on frequency ω and sideband on frequencies $\omega \pm \omega_M$). These fields can induce a Raman transition between the two Bloch bands of a BEC $s = 0$ and $s \neq 0$. A resonant field via a two - photon (Raman) transition changes the population of the lowest Bloch band $s = 0$ of a BEC. In resonant case the system is excited by an off - resonant Raman (two - photon) driving with the Rabi frequency Ω_R . In an amplitude - modulated optical lattice two standing laser waves of frequencies ω and $\omega - \omega_M$ ($\omega_M \ll \omega$) drive an atom in a Raman scheme. The effective two - photon Rabi frequency is given in terms of the single - photon Rabi frequencies Ω and Ω' of the fields on the main ω and sideband $\omega' = \omega - \omega_M$ frequencies as $\Omega_R = \frac{\Omega \cdot \Omega'}{2\Delta}$. For the amplitude - modulated laser beams with the modulation strength ν we have for standing waves $\Omega_R(\mathbf{x}) = \frac{\nu \Omega^2(\mathbf{x})}{2\Delta} = \frac{\nu V_0(\mathbf{x})}{\hbar}$, here Δ is the far - detuning of the laser beams from the atomic resonance (the closest neighboring optical dipole transition), and $V_0(\mathbf{x}) = \sum_{j=1}^3 V_{jo} \cos^2 kx_j$ is the optical lattice potential with wave vectors $k = \frac{2\pi}{\lambda}$ and λ the wavelength of the laser beams on the main frequency, and we are omitted the terms proportional to the small parameter $\Delta k \cdot x = \frac{\omega_M}{c} x \ll 1$, where $\Delta k \equiv k' - k$, for $x < \lambda \sim 10^{-4} \text{cm}$, $\omega_M \preceq 10^7 \text{s}^{-1}$, this parameter is about 10^{-7} . The eigenstates of a BEC are Bloch waves, and an appropriate superposition of Bloch states yields a set of Wannier functions which are localized on the individual lattice sites. Without a resonant driving the wave functions in a tight binding model can be represented [12] as $w_s(\mathbf{x}) = Z_s \Psi_s(\mathbf{x})$, where $\Psi_s(\mathbf{x})$ is the wave function of the oscillator in a state s , and Z_s is a normalization factor. However, for a resonant driving the wave functions are

$w_s^{res}(\mathbf{x}) = \varphi(\mathbf{x})w_s(\mathbf{x})$, where the function $\varphi(\mathbf{x})$ is the solution of the Bloch equation of the standard Rabi problem. We consider a steady state driving regime, in which the function $\varphi(\mathbf{x})$ is determined on the parameters of the field (the Rabi frequency and the detuning from resonance $\delta = \omega_M - (\omega_b - \omega_a)$). In order to avoid a sufficient population of the excited Bloch band $s \neq 0$, we are assume $\Omega_R < \delta$, then $\varphi^2(\mathbf{x}) \simeq 1 - \frac{1}{2}(\frac{\Omega_R(\mathbf{x})}{\delta})^2$. Using the wave functions w_s^{res} , in which the size of the ground state oscillator wave function l_0 in a lattice is much less than the lattice period $d = \lambda/2$, and following Ref. [12] we find the effective scattering length a_{eff}^{res} modified by the resonantly amplitude modulation

$$\frac{a_{eff}^{res} - a_{eff}}{a_{eff}} \simeq -(\frac{\nu V_0}{h\delta})^2. \quad (3)$$

As a result, we can manipulate the effective scattering length in a BEC in an resonantly amplitude - modulated optical lattice. This tuning is determined by the modulation strength ν and the detuning δ . By varying the parameter ν/δ , we can tune a_{eff}^{res} to a given value.

A nonresonant case.

Second is a nonresonant mechanism for which we will use a direct time - dependent description. Note that such potential is created by interfering pair of amplitude - modulated laser beams with the modulation strength $\nu/2$ of the electric field amplitude, and was omitted the quadratic term for the small parameter $\nu^2/4$. A nonresonant low - frequency mechanism of manipulation is in operation for a quasistatic regime $\omega_M < 2\pi\tau_{noneq}^{-1}$, therefore, if V_0 corresponds to a geometric resonance at $l_* = |a|$ for attractive interaction then as the time runs, the system goes from an attractive interaction to a repulsive interaction and back again with frequency ω_M . Therefore, it is allow to investigate the nonequilibrium dynamics in a BEC. In a nonresonant case $\Omega_R/\delta \rightarrow 1$ then $\varphi(\mathbf{x}) = 1$ and we have a time - dependent WKB tunneling exponent $D_0(t)$. The WKB tunneling exponent is a nonlinear function on the strength of the periodic potential V_0 which is proportional to the laser intensity . If we have a time - dependent lattice laser intensity, then, although $\overline{V_0(t)} = V_0$, due to the nonlinearity of a WKB tunneling exponent a time - averaged $\overline{D_0(t)}$ will be different from a WKB tunneling exponent D_0 without a modulation. For a small ν we have the

time - averaged $\overline{D_0(t)} = D_0 + \Delta D_0$, where the bar stands for a time average and ΔD_0 is proportional to the time - averaged $\nu^2 \overline{\cos^2 \omega_M t}$ (the time - averaged of the modulation harmonic oscillation to the lowest even order power). For a low - frequency nonresonant manipulation we will use a time - dependent potential (Eq.(2)), and there could be a modulation frequency ω_M low enough for quasistatic behavior. Using the WKB tunneling exponent $D_0 \simeq \exp(-\frac{\pi d}{\hbar} \sqrt{2mV_0})$ and using $V_0(t)$ (Eq.(2)), we have a time - dependent length scale $l_*^{non}(t)$ and, correspondingly, a time - dependent effective scattering length $a_{eff}^{non}(t)$. Calculating the time - average $\overline{D_0(t)}$ by expanding in a series about the small parameter $2mV_0\pi^2 d^2/\hbar^2 < 1$, we find the time - average length scale $\overline{l_*^{non}}$, which is modified by the nonresonant amplitude modulation

$$\frac{\overline{l_*^{non}} - l_*}{l_*} \simeq \frac{\nu^2}{4} \frac{2mV_0\pi^2 d^2}{\hbar^2}. \quad (4)$$

If $\frac{\nu^2}{4} \frac{2mV_0\pi^2 d^2}{\hbar^2}$ is not small, then we may expand the expression for the time - dependent $D_0(t)$ using Bessel functions $I_n(x)$ and the functions $\cos^n \omega_M t$. In this case instead the parameter $\frac{\nu^2}{4} \frac{2mV_0\pi^2 d^2}{\hbar^2}$ we have $I_2(\nu \frac{\pi d}{\hbar} \sqrt{2mV_0})$, because, usually $\frac{\pi d}{\hbar} \sqrt{2mV_0} \gg 1$ and for $\nu < 0.4$ among $I_n(x < 2)$ the largest is $I_2(x)$.

In summary, the effective scattering length in a BEC in an optical lattice can be manipulated by an amplitude - modulated manner. The effect of the modulation is to modulate the optical lattice potential so that it becomes a time - dependent optical lattice potential depth and, correspondingly, the effective scattering length becomes time - dependent. There are two mechanisms of the manipulation. One, a resonant mechanism is in operation for the modulation frequency closed to the excited internal state of a BEC, and two standing laser waves of main and sideband frequencies of an amplitude - modulated optical lattice induce a Raman transition. Due to resonant Raman driving the population of the lowest Bloch band $s = 0$ of a BEC changes, therefore, the effective scattering length is also changed. As a result, the effective scattering length is other than without a modulation, this tuning is determined by the square of the modulation strength and the square of the detuning. Second, a nonresonant low - frequency mechanism, for which we use a direct time - depen-

dent description. Due to the nonlinear dependence of the WKB tunneling amplitude on the optical lattice depth for the harmonic modulation of the optical lattice depth the time - averaged tunneling exponent $\overline{D_0(t)}$ is different from D_0 without a modulation. The proposed mechanism of manipulating the effective scattering length makes it possible to tune it to a given value. As a result, although a time - averaged optical lattice potential with a harmonic modulation is equal to an optical lattice potential without a modulation, the effective scattering length in a BEC confined in an amplitude - modulated optical lattice can be manipulated by the modulation strength. In conclusion, the nonequilibrium dynamics of a BEC can be investigated by manipulating the effective scattering length of a BEC in an amplitude - modulated optical lattice.

I thank A.J.Freeman for encouragement.

REFERENCES

- [1] M. H. Anderson *et al.*, Science **269**, 198 (1995).
- [2] C. C. Bradley *et al.*, Phys. Rev. Lett. **75**, 1687 (1995).
- [3] K. B. Davis *et al.*, Phys. Rev. Lett. **75**, 3969 (1995).
- [4] C. Orzel *et al.*, Science **291**, 2386 (2001).
- [5] M. Greiner *et al.*, Nature (London) **415**, 39 (2002).
- [6] F. S. Cataliotti *et al.*, Science **293**, 843 (2001).
- [7] O. Morsch *et al.*, Phys. Rev. Lett. **87**, 140402 (2001).
- [8] M. Olshanii, Phys. Rev. Lett. **81**, 938 (1998).
- [9] D. S. Petrov, G. V. Shlyapnikov, and J. T. M. Walraven, Phys. Rev. Lett. **85**, 3745 (2000).
- [10] D. S. Petrov, M. Holzmann, and G. V. Shlyapnikov, Phys. Rev. Lett. **84**, 2551 (2000).
- [11] D. S. Petrov and G. V. Shlyapnikov, Phys. Rev. A **64**, 012706 (2001).
- [12] P. O. Fedichev, M. J. Bijlsma, and P. Zoller, Phys. Rev. Lett. **92**, 080401 (2004).
- [13] A. Polkovnikov, S. Sachdev, and S. M. Girvin, Phys. Rev. A **66**, 053607 (2001).
- [14] K. Berg - Sørensen and K. Mølmer, Phys. Rev. A **58**, 1480 (1998).
- [15] D. Diakonov *et al.*, Phys. Rev. A **66**, 013604 (2002).